

Four new species of *Erioscyphella* (Leotiomyces, Helotiales) from southwestern China

Le Luo^{1,2,3,4}, Kandawatte W. Thilini Chethana^{2,3}, Qi Zhao¹, Hong-Li Su^{1,2,3}, Cui-Jin-Yi Li^{1,2,3},
Vinodhini Thiyagaraja¹, Fatimah Al-Otibi⁵, Kevin D. Hyde^{1,2,5}

¹ Key Laboratory of Phytochemistry and Natural Medicines, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, Yunnan, China

² Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

³ School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

⁴ Guiyang Institute of Humanities and Technology, Guiyang 550025, Guizhou, China

⁵ Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 22452, Riyadh 11495, Saudi Arabia

Corresponding authors: Kandawatte W. Thilini Chethana (kandawatte.thi@mfu.ac.th); Kevin D. Hyde (kdhyde3@gmail.com)



This article is part of:

**Exploring the Hidden Fungal Diversity:
Biodiversity, Taxonomy, and Phylogeny of
Saprobic Fungi**

Edited by Samantha C. Karunarathna,
Danushka Sandaruwan Tennakoon, Ajay
Kumar Gautam

Academic editor:

Danushka Sandaruwan Tennakoon

Received: 5 October 2024

Accepted: 26 November 2024

Published: 14 February 2025

Citation: Luo L, Thilini Chethana KW, Zhao Q, Su H-L, Li C-J-Y, Thiyagaraja V, Al-Otibi F, Hyde KD (2025) Four new species of *Erioscyphella* (Leotiomyces, Helotiales) from southwestern China. MycoKeys 114: 29–48. <https://doi.org/10.3897/mycokeys.114.138647>

Copyright: © Le Luo et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Abstract

Erioscyphella is found across various regions and is part of the family Lachnaceae (Helotiales). It is distinguished by its white to orange disc-shaped apothecia, white to brown receptacles, and granulated hairs that contain amorphous or resinous material. These hairs lack swelling apices and crystals. Additionally, this genus is unique for its long ascospores. In the present study, we collected eight specimens from southwestern China. Morphological and phylogenetic analyses based on the combined LSU, ITS, mtSSU and *RPB2* dataset showed that our specimens represent four new species of *Erioscyphella*, including *E. ailaoensis*, *E. baimana*, *E. gelangheica* and *E. tengyueica*. Here, we provide complete morphological descriptions with illustrations and sequence data essential for future taxonomic and evolutionary research.

Key words: 4 novel species, Lachnaceae, morphology, phylogeny, taxonomy

Introduction

The monophyletic genus *Erioscyphella* belongs to the family Lachnaceae (Dennis 1954; Spooner 1987; Cantrell and Haines 1997; Guatimosim et al. 2016; Tochiara and Hosoya 2022; Wijayawardene et al. 2020, 2022; Hyde et al. 2024) and includes 22 records (Index Fungorum 2024). The original description of *Erioscyphella* by Kirschstein (1939), which lacked typification, was inaccurately characterized based on traits that lack taxonomic significance, such as filiform, colored, and pigmented ascospores, as well as lanceolate paraphyses (Korf 1978; Perić and Baral 2014). It was not until Haines (1984) selected *E. longispora*, under which *Peziza abnormis* was later synonymized, as the lectotype of *Erioscyphella* that a more accurate characterization was established. These features are now recognized as insufficient for distinguishing this genus from related taxa. This genus was initially confused with the closely related genus *Lachnum*, but detailed morphological and molecular phylogenetic studies based on LSU, ITS, mtSSU and

RPB2 data have since clarified their distinct characteristics and proposed a new concept based on the examination of Japanese materials (Tochihara and Hosoya 2022). *Erioscyphella* is characterized by scattered and cupulate apothecia on leaves or bamboo sheaths, straight or irregularly curved, septate and granulated hairs, mostly covered by apical amorphous materials or resinous material, lanceolate or filiform paraphyses, 8-spored asci with an amyloid apical pore and fusiform to long needle-like ascospores (Kirschstein 1939; Perić and Baral 2014; Tochihara and Hosoya 2022). There are no reports of asexual morph in this genus.

The species of *Erioscyphella* are primarily distributed in temperate and tropical regions, commonly inhabiting decaying wood and plant debris. *Erioscyphella* species are distributed mainly in China and a few other countries; *E. curvispora* is described from Montenegro (Perić and Baral 2014), *E. griseibambusicola*, *E. latispora*, *E. lunata*, *E. lushanensis* and *E. subinsulae* are collected from China (Guatimosim et al. 2016; Tello and Baral 2016; Li et al. 2022; Su et al. 2023); *E. euterpes* collected from Puerto Rico (Guatimosim et al. 2016). In contrast to the above-mentioned species that show a narrower distribution, *E. abnormis* shows a worldwide distribution, especially endemic in tropical regions (Tello and Baral 2016). They play a crucial role in breaking down organic matter, thereby aiding in nutrient cycling within their ecosystems. The ecological role of *Erioscyphella* as decomposers underscores their importance in forest ecosystems (Tochihara and Hosoya 2022; Niego et al. 2023a, 2023b). Ongoing research aims to better define species boundaries and elucidate the phylogenetic relationships within the genus through morphological studies and DNA sequencing. This work aids in the precise classification and discovery of new species.

During the investigation of Leotiomyces in southwest China (Li et al. 2022, 2024a, 2024b; Su et al. 2022, 2023; Luo et al. 2024; Thiyagaraja et al. 2024), eight collections of *Erioscyphella* were obtained. We used morphological and phylogenetic analyses based on LSU, ITS, mtSSU and *RPB2* data to confirm that these eight collections differ from all known species of *Erioscyphella*. We introduce four species to accommodate these collections. Here, we provide complete morphologies, illustrations, and their phylogenetic relationships for future taxonomic and evolutionary studies.

Material and methods

Specimen collection and morphological examination

We collected eight specimens from southwest China. All samples were collected from highly humid, natural broadleaf forests and protected areas with minimal human access. Altitudes were determined by the GPS device. The fruiting bodies were discovered on the surface of extremely wet, decaying wood litter. The samples were dehydrated in a dehydrator at a temperature range of 25–30 °C. After studying the morphology of the specimens and getting their genomic DNA, they were deposited at the Cryptogamic Herbarium of the Kunming Institute of Botany, Chinese Academy of Sciences (KUN-HKAS). Facesoffungi and Index Fungorum numbers were obtained as in Jayasiri et

al. (2015) and Index Fungorum (2024), respectively. All the species identifications followed Chethana et al. (2021). The morphological descriptions were submitted to the Greater Mekong Subregion database (Chaiwan et al. 2021). The dried specimens were examined with a stereomicroscope (C-PSN, Nikon, Japan) and were captured with a digital camera (Canon EOS 70D, Japan) connected to the stereomicroscope. Free-hand sections of the dried specimens were mounted in a drop of water for observing microscopic characteristics, such as apothecia, exciple, paraphyses, asci and ascospores, using a Nikon compound microscope (Nikon, Japan) equipped with a DS-Ri2 camera. In addition, the sections were pretreated with Melzer's reagent for the Iodine test (MLZ) (Tochihara and Hosoya 2022). Microstructures were measured using the Tarosoft (R) Image Frame Work program v.0.97 (Tarosoft, Thailand). The obtained measurements were presented in the format of (a–) b–c(–d), where 'a' represented the minimum value, 'd' represented the maximum value, and the range 'b–c' reflected the 90% confidence interval. The $\bar{x}$ indicated the average value of measurements. Ascospore measurements were given as [n/m/p], indicating that the n number of ascospores were measured from m ascomata of the p number of collections. Images used for figures were processed with Adobe Photoshop CS6 Extended version 13.0 × 64 (Adobe Systems, USA).

DNA extraction, PCR amplifications and sequencing

Genomic DNA was extracted from the dried apothecia (around 50–100 mg) using a TSP101 DNA extraction kit (TSINGKE, China). Following the latest studies (Tochihara and Hosoya 2022; Su et al. 2023; Luo et al. 2024), LSU, ITS, mtSSU and *RPB2* were used for PCR amplification, using the primers LR0R/LR5 (Vilgalys and Hester 1990), ITS1F/ITS4 (White et al. 1990; Gardes and Bruns 1993), mrSSU1/mrSSU3R (Zoller et al. 1999) and *fRPB2*-5F/*fRPB2*-7cR (Liu et al. 1999), respectively. For LSU, ITS, mtSSU and *RPB2*, the total volume of PCR amplifications was 25 µL, comprising 12.5 µL 2 × PCR G013 Taq MasterMix with Dye (Applied Biological Materials, Canada), 1 µL of each primer (10 µM), 2 µL genomic DNA, and 8.5 µL of sterilized, distilled water. Amplifications of LSU, ITS and *RPB2* were conducted under the following conditions: pre-denaturation at 95 °C for 5 min, followed by 35 cycles of denaturation at 95 °C for 20 sec, annealing at 56 °C (LSU)/53 °C (ITS and *RPB2*) for 10 sec, elongation at 72 °C for 20 sec, and final elongation at 72 °C for 7 min. For mtSSU, the total volume of PCR amplifications was 25 µL, which comprised 21 µL 1 × PCR TSE101 Mix (TSINGKE, China), 1 µL of each primer (10 µM), and 2 µL genomic DNA. Amplifications of mtSSU were conducted under the following conditions: initial denaturation at 98 °C for 3 min, followed by 40 cycles of denaturation at 98 °C for 1 min, annealing at 52 °C for 1 min, elongation at 72 °C for 1 min, and final elongation at 72 °C for 10 min. Gel electrophoresis with 1% TAE and TSJ003 GoldView nucleic acid dye (TSINGKE, China) was used to confirm the obtained PCR products. Finally, the PCR products were sequenced at the Tsingke Biotechnology Co., Ltd., Kunming, China. Newly produced sequences were deposited in the GenBank and the accession numbers were given in Table 1.

Table 1. Taxa included in the phylogenetic analyses and the GenBank accession numbers of LSU, ITS, mtSSU and *RPB2* sequences.

Species	Strain	Gene accession No.				References
		ITS	LSU	mtSSU	<i>RPB2</i>	
<i>Capitotricha bicolor</i>	TNS-F-65670	LC424834	LC424942	LC533244	LC425011	Tochihara and Hosoya (2022)
<i>Capitotricha rubi</i>	TNS-F-65752	LC438560	LC438573	LC533243	LC440395	Tochihara and Hosoya (2022)
<i>Erioscyphella abnormis</i>	TNS-F-16609	AB705234	LC533175	LC533256	LC533184	Tochihara and Hosoya (2022)
<i>Erioscyphella abnormis</i>	TNSF38452	LC669457	LC533171	LC533262	LC533210	Tochihara and Hosoya (2022)
<i>Erioscyphella abnormis</i>	TNS-F-80478	LC424837	LC424949	LC533283	-	Tochihara and Hosoya (2019)
<i>Erioscyphella abnormis</i>	TNS-F-46841	LC669474	LC533170	LC533279	LC533209	Tochihara and Hosoya (2022)
<i>Erioscyphella ailaoensis</i>	HKAS135686^(T)	PQ349783	PQ349775	PQ358800	PQ424108	This study
<i>Erioscyphella ailaoensis</i>	HKAS135687	PQ349784	PQ349776	PQ358801	PQ424109	This study
<i>Erioscyphella alba</i>	MFLU16-0614 ^(T)	MK584965	MK591990	-	-	Ekanayaka et al. (2019)
<i>Erioscyphella aseptata</i>	MFLU16-0590 ^(T)	MK584957	MK591986	-	MK388223	Ekanayaka et al. (2019)
<i>Erioscyphella baimana</i>	HKAS135697^(T)	PQ349785	PQ349777	PQ358802	PQ424110	This study
<i>Erioscyphella baimana</i>	HKAS135696	PQ349786	PQ349778	PQ358803	PQ424111	This study
<i>Erioscyphella boninensis</i>	TNS-F-26520 ^(T)	NR185389	LC533151	LC533254	LC533196	Tochihara and Hosoya (2022)
<i>Erioscyphella brasiliensis</i>	MFLU16-0577b	MK584967	MK591993	-	-	Ekanayaka et al. (2019)
<i>Erioscyphella brasiliensis</i>	TNS-F-46419	LC669456	LC533133	LC533278	LC549672	Tochihara and Hosoya (2022)
<i>Erioscyphella curvispora</i>	KL 381 ^(T)	MH190414	MH190415	-	-	Perić and Baral (2014)
<i>Erioscyphella euterpes</i>	PR 147	U58640	-	-	-	Cantrell and Haines (1997)
<i>Erioscyphella fusiforme</i>	MFLU15-0230 ^(T)	MK584948	MK591975	-	MK614728	Ekanayaka et al. (2019)
<i>Erioscyphella gelangheica</i>	HKAS135689^(T)	PQ349787	PQ349779	PQ358804	-	This study
<i>Erioscyphella gelangheica</i>	HKAS135695	PQ349788	PQ349780	PQ358805	-	This study
<i>Erioscyphella griseibambusicola</i>	HKAS124657	OP451797	OP451791	OP451844	OP432252	Su et al. (2023)
<i>Erioscyphella griseibambusicola</i>	HKAS124656 ^(T)	OP451796	OP451790	OP451843	OP432251	Su et al. (2023)
<i>Erioscyphella hainanensis</i>	TNS-F-35056	LC669465	LC533169	LC533275	LC533206	Tochihara and Hosoya (2022)
<i>Erioscyphella hainanensis</i>	TNS-F-35049	LC669452	LC533168	LC533274	LC533205	Tochihara and Hosoya (2022)
<i>Erioscyphella insulae</i>	TNS-F-26500	LC669448	LC533149	LC533252	LC533194	Tochihara and Hosoya (2022)
<i>Erioscyphella insulae</i>	TNS-F-39720 ^(T)	LC669451	LC533177	LC533261	LC533207	Tochihara and Hosoya (2022)
<i>Erioscyphella latispora</i>	HKAS124391	OP113849	OP113850	-	OP715727	Li et al. (2022)
<i>Erioscyphella latispora</i>	HKAS124389 ^(T)	OP310823	OP113844	-	OP715728	Li et al. (2022)
<i>Erioscyphella lunata</i>	JA-CUSSTA 8292	KX501132	KX501133	-	-	Tello and Baral (2016)
<i>Erioscyphella lushanensis</i>	HMAS81575	JF937582	-	-	-	Zhao and Zhuang (2011)
<i>Erioscyphella otanii</i>	TNS-F-81472 ^(T)	NR185393	LC533179	LC533286	LC533226	Tochihara and Hosoya (2022)
<i>Erioscyphella papillaris</i>	TNS-F-81272 ^(T)	NR185391	LC533161	LC533285	LC533204	Tochihara and Hosoya (2022)
<i>Erioscyphella paralushanensis</i>	TNS-F-61920 ^(T)	NR185390	LC533141	LC533267	LC533220	Tochihara and Hosoya (2022)
<i>Erioscyphella sasibrevispora</i>	TNS-F-80399	LC669470	LC533173	LC533268	LC533216	Tochihara and Hosoya (2022)
<i>Erioscyphella sasibrevispora</i>	TNS-F-81401 ^(T)	LC669472	LC533174	LC533269	LC533217	Tochihara and Hosoya (2022)
<i>Erioscyphella sclerotii</i>	TNS-F-26492	LC669438	LC533152	LC533255	LC533197	Tochihara and Hosoya (2022)
<i>Erioscyphella sclerotii</i>	TNS-F-38480	LC669458	LC533134	LC533263	LC549673	Tochihara and Hosoya (2022)
<i>Erioscyphella sclerotii</i>	MFLU 16-0569	MK584951	MK591980	-	-	Ekanayaka et al. (2019)
<i>Erioscyphella sclerotii</i>	MFLU 18-0688	MK584969	MK591995	-	-	Ekanayaka et al. (2019)
<i>Erioscyphella sinensis</i>	TNS-F-32161	LC669449	LC533167	LC533273	LC533219	Tochihara and Hosoya (2022)
<i>Erioscyphella sinensis</i>	TNS-F-16838	AB481280	LC533164	LC533235	AB481364	Tochihara and Hosoya (2022)
<i>Erioscyphella subinsulae</i>	HKAS 124659	OP451799	OP451793	OP451846	OP432254	Su et al. (2023)
<i>Erioscyphella subinsulae</i>	HKAS 124660	OP451800	OP451794	OP451847	OP432255	Su et al. (2023)
<i>Erioscyphella subinsulae</i>	HKAS 124661	OP451801	OP451795	OP451848	OP432256	Su et al. (2023)
<i>Erioscyphella subinsulae</i>	HKAS 124658 ^(T)	OP451798	OP451792	OP451845	OP432253	Su et al. (2023)
<i>Erioscyphella tengyueica</i>	HKAS135688^(T)	PQ349789	PQ349781	PQ358806	-	This study
<i>Erioscyphella tengyueica</i>	HKAS135693	PQ349790	PQ349782	PQ358807	-	This study
<i>Lachnellula calyciformis</i>	TNS-F-81248	LC438561	LC438574	LC533247	LC438590	Tochihara and Hosoya (2022)
<i>Lachnellula suecica</i>	TNS-F-16529	AB481248	LC424944	LC533231	AB481341	Tochihara and Hosoya (2022)
<i>Neodasyscypha cerina</i>	TNS-F-65625	LC424836	LC424948	LC533242	LC425013	Tochihara and Hosoya (2022)

*Names in indicate the specimens from the current study. Names with ^(T) indicate type specimens and - denotes unavailable data in the GenBank.

Phylogenetic analyses

New DNA sequences generated from forward and reverse primers were assembled using BioEdit v.7.2.5 (Hall 1999) to obtain consensus sequences. The concatenated sequences were used to search for the closer relatives in the NCBI (Johnson et al. 2008). According to the close relatives and recent studies, the newly generated sequences and some published sequences were used for the phylogenetic analyses (Table 1). *Capitotricha bicolor* (TNS-F-65670), *C. rubi* (TNS-F-65752), *Lachnellula calyciformis* (TNS-F-81248), *L. suecica* (TNS-F-16529) and *Neodasyscypha cerina* (TNS-F-65625) were used as the outgroup taxa. The phylogenetic analysis was conducted based on the datasets including reference DNA sequences and newly generated DNA sequences using OFPT (Zeng et al. 2023) with the following protocol. Datasets of each gene region were first independently aligned with ‘auto’ strategy (based on data size) by MAFFT (Kato and Standley 2013) and trimmed with ‘gappyout’ method (based on gaps’ distribution) by TrimAl (Capella-Gutiérrez et al. 2009). The best-fit nucleotide substitution model for each dataset was then selected based on the Bayesian information criterion (BIC) from twenty-two common DNA substitution models with rate heterogeneity by ModelFinder (Kalyana-moorthy et al. 2017). Afterwards, all datasets were concatenated with partition information for the subsequent phylogenetic analyses. Maximum likelihood with 1000 replicates was performed using ultrafast bootstrap approximation (Hoang et al. 2018) with SH-like approximate likelihood ratio test (SH-aLRT) (Guindon et al. 2010) in IQ-TREE (Nguyen et al. 2015). The consensus tree was summarized based on the extended majority rule. Bayesian inference (BI) analyses were run in the CIPRES Science Gateway v.3.3 (Miller et al. 2010). The best-fit nucleotide substitution models were determined using jModelTest2 on XSEDE (2.1.6). The BI was performed in MrBayes on XSEDE v. 3.2.7a (Ronquist et al. 2012), with four simultaneous Markov chain Monte Carlo (MCMC) chains and four runs for 3,000,000 generations, with trees sampled at each 300th generation. The first 25% of trees were discarded as burn-in, and BI posterior probabilities (PP) were conducted from the remaining trees. The consensus phylograms were visualized on FigTree v. 1.4.4, and edited with Adobe Illustrator CC 2019, Adobe Systems (USA). Decisions as to whether species are new followed the polyphasic approach as recommended by Chethana et al. (2021) and Maharachchikumbura et al. (2021).

Results

Phylogenetic analysis

The phylogenetic analyses were based on 45 *Erioscyphella* taxa, including *C. rubi* (TNS-F-65752), *C. bicolor* (TNS-F-65670), *L. calyciformis* (TNS-F-81248), *L. suecica* (TNS-F-16529) and *N. cerina* (TNS-F- 65625) as the outgroup taxa. The alignment comprised 4 partitions and 3944 total sites (ITS: 758 bp; LSU: 1105 bp; mtSSU: 947 bp; *RPB2*: 1134 bp), with 14.359% gaps and completely undetermined characters. The ML tree has the same topology as the BI tree. The best ML tree with a final optimization likelihood of -25316.078809 is displayed in Fig. 1. In the BI analyses, the final average standard deviation of split




Figure 1. The Maximum Likelihood tree based on the combined LSU, ITS, mtSSU and *RPB2* sequence data for *Erioscyphella*. *Capitotricha bicolor* (TNS-F-65670), *C. rubi* (TNS-F-65752), *Lachnellula calyciformis* (TNS-F-81248), *L. suecica* (TNS-F-16529) and *Neodasyscypha cerina* (TNS-F-65625) are used as the outgroup taxa. The MLBP $\geq 70\%$ and BPP ≥ 0.90 are shown at the nodes as MLBP/BPP. MLBS $< 70\%$ and BPP < 0.90 are expressed as a hyphen (“-”). Names with (T) indicate type specimens. Names in red indicate new species.

frequencies was 0.007666, which revealed convergence. In the multi-gene phylogenetic tree based on the combined ITS, LSU, mtSSU and *RPB2* dataset, all taxa of *Erioscyphella* clustered together. *Erioscyphella ailaoensis* was sister to the clade comprising *E. sclerotii* and *E. abnormis*, with 73% maximum likelihood bootstrap (MLBP) and 0.89 Bayesian posterior probabilities (BPP) support. *Erioscyphella gelangheica* was sister to the clade comprising *E. sclerotii* and *E. abnormis*, *E. ailaoensis*, *E. brasiliensis* and *E. aseptata* with 78% MLBP and 0.88 BPP support. *Erioscyphella baimana* was sister to *E. latispora* with 99% MLBP and 1.00 BPP support. Besides, *E. tengyueica* was sister to *E. papillaris* with 91% MLBP and 0.98 BPP. The phylogenetic result showed that *Erioscyphella* species clustered together, similar to previously published studies (Su et al. 2023).

Taxonomy

***Erioscyphella ailaoensis* L. Luo, K.D. Hyde & H.L. Su, sp. nov.**

Index Fungorum: IF902529

Facesoffungi Number: FoF16389

Fig. 2

Etymology. The epithet “*ailaoensis*” refers to the collection site, Ailao Mountain, where the holotype specimen was collected.

Holotype. HKAS135686.

Description. **Saprobic** on the dead bark. **Sexual morph:** **Apothecia** scattered to partly gregarious, superficial, 1–2.4 mm in diameter, 0.4–1.4 mm high when dry, discoid to cupulate, shortly stipitate, externally covered with short, white to brown hairs. **Disc** concave, surface slightly rough, yellow to brown. Margin flat to slightly involute, pale yellow, covered with yellow to pale brown hairs. **Receptacle** discoid to cupulate, yellow to pale brown, clothed entirely with short, yellow to pale brown hairs. **Stipe** 0.2–1 mm in diameter, 0.2–0.6 mm long when dry, cylindrical, solitary, yellow to pale brown, clothed with yellow to pale brown hairs. **Hairs** 22–92 × 3.0–4.3 µm ($\bar{x}$ = 51 × 3.7 µm, n = 30), clavate to cylindrical, straight to slightly curved, septate, hyaline, thick-walled, covered with hyaline granules, obtuse apex, apical amorphous or resinous material. **Hymenium** 120–230 µm ($\bar{x}$ = 153 µm, n = 12), concave, surface slightly rough, light yellowish brown in dry. **Medullary excipulum** 40–90 µm ($\bar{x}$ = 58 µm, n = 20), thin, hyaline to light yellow, thin-walled, smooth cells of **textura oblita**, 2.1–5.9 µm ($\bar{x}$ = 3.7 µm, n = 50) in diameter. **Ectal excipulum** 55–80 µm ($\bar{x}$ = 65 µm, n = 20), thin-walled, smooth, light yellowish cells of **textura prismatica** to **globulosa**, 2–4.9 µm ($\bar{x}$ = 3.0 µm, n = 60) in diameter. **Paraphyses** 100–138 × 1.6–3.9 µm ($\bar{x}$ = 117 × 2.8 µm, n = 25), longer than asci, filiform, straight to slightly curved, aseptate, hyaline, thin-walled, rough, with slightly acute apex. **Asci** 85–143 × (4.5–) 5.5–9.0(–9.5) µm ($\bar{x}$ = 100 × 7.3 µm, n = 34), 8-spored, unitunicate, overlapping fascicles, clavate, straight to slightly curved, inoperculate, hyaline, apically thickened wall, laterally relatively thin, slightly smooth, with an apical, non-amyloid pore and tapered ends, J- in MLZ. **Ascospores** (50/14/2) (43.0–)45.5–97(–101.0) × 1.4–2.4(–2.6) µm, ($\bar{x}$ = 68 × 1.9 µm), fascicled, filiform, multi-septate, thin-walled, hyaline, rough with taper, obtuse ends, without oil guttules, hyaline, slightly smooth. **Asexual morph:** Not observed.



Figure 2. *Erioscyphella ailaoensis* (HKAS135686, holotype) **a–c** dried ascomata on the bark **d** vertical section of an ascoma **e** excipulum **f** hairs **g** paraphyses **h–k** asci (**j, k** asci in MLZ) **l–o** ascospores (**l–n** ascospores in MLZ). Scale bars: 100 μ m (**d**); 50 μ m (**e–k**); 30 μ m (**l–o**).

Material examined. CHINA • Yunnan Province, Puer City, Jingdong County, Ailao Mountain, altitude 2478 m, on the decayed unidentified bark, 8 June 2022, Hongli Su, SU872 (HKAS135686, **holotype**); CHINA • Xizang Province, Shigatse City, altitude 1774 m, on the decayed unidentified twig, 6 July 2022, Hongli Su, SU1423 (HKAS135687, **paratype**).

Notes. Our specimens, HKAS135686 and HKAS135687, were grouped as a distinct clade, separated from the clade comprising *E. abnormis* and *E. sclerotii* by 73% MLBS and 0.89 BIPP (Fig. 1). The new species exhibited morphological

differences from *E. abnormis* and *E. sclerotii* by having J- apical pores, whereas the latter species have apical pores that are J- in MLZ. In contrast to the septate paraphyses of *E. abnormis*, *E. ailaoensis* has aseptate paraphyses (Han et al. 2021). Furthermore, asci and ascospores of *E. ailaoensis* are longer than those of *E. sclerotii* (Nagao 1996; Perić and Baral 2014). Therefore, *E. ailaoensis* is introduced here as a new species.

***Erioscyphella baimana* L. Luo, K.D. Hyde & Q. Zhao, sp. nov.**

Index Fungorum: IF902530

Facesoffungi Number: FoF16391

Fig. 3

Etymology. The epithet “*baimana*” refers to the collection site, Baima Mountain, where the holotype specimen was collected.

Holotype. HKAS 135697.

Description. *Saprobic* on dead twigs. **Sexual morph: Apothecia** superficial, gregarious, 0.3–1.1 mm in diameter, 0.3–1.4 mm high when dry, discoid to cupulate, long stipitate, externally covered with short, white hairs. **Disc** concave, surface slightly smooth, yellow. **Margin** flat to slightly involute, white, covered with white hairs. **Receptacle** cupulate to discoid, white, covered entirely with short, white hairs. **Stipe** 0.2–0.6 mm in diameter, 0.3–1.1 mm long when dry, cylindrical, solitary, white, clothed with white hairs. **Hairs** 30–120 × 2.8–4.7 µm ($\bar{x}$ = 74 × 3.7 µm, n = 30), clavate to cylindrical, straight to slightly curved, aseptate, hyaline, thin-walled, covered with fine granules, obtuse apex, lacks apical amorphous. **Hymenium** 165–230 µm ($\bar{x}$ = 195 µm, n = 12), concave, surface slightly smooth, yellow in dry. **Medullary excipulum** 35–120 µm ($\bar{x}$ = 70 µm, n = 18), thin, hyaline, thin-walled cells of **textura porrecta**, 1.3–3.8 µm ($\bar{x}$ = 2.5 µm, n = 50) in diameter. **Ectal excipulum** 40–120 µm ($\bar{x}$ = 68 µm, n = 18) thin, thin-walled, smooth, light yellowish cells of **textura porrecta** to **oblita**, 1.9–6.3 µm ($\bar{x}$ = 4.1 µm, n = 60) in diameter. **Paraphyses** 95–170 × 1.2–2.5 µm ($\bar{x}$ = 140 × 1.6 µm, n = 25), longer than asci, filiform, straight to slightly curved, aseptate, hyaline, light smooth, with slightly obtuse apex. **Asci** 100–152 × 3.6–9.5 µm ($\bar{x}$ = 123 × 7.2 µm, n = 34), 8-spored, unitunicate, clavate, straight to slightly curved, inoperculate, hyaline, slightly smooth, with an apical, amyloid pore and rounded ends, J+ in MLZ, tapered long stipitate base. **Ascospores** (85/7/2) (29.5–)30.5–37.5(–40.0) × 2.2–5.0 (–5.5) µm ($\bar{x}$ = 32.9 × 4.1 µm, n = 85), biserial, fusoid-clavate with blunt ends, fusiform, 1–3-septate, thin-walled, hyaline, slightly smooth, tapering towards the obtuse ends, with longitudinal striations, without oil guttules. **Asexual morph:** Not observed.

Material examined. CHINA • Yunnan Province, Diqing City, Deqin County, Baima Mountain, altitude 3485 m, on the decayed unidentified twig, 21 July 2022, Le Luo, Ly7 (HKAS 135697, **holotype**); • *ibid.*, Le Luo, Ly26 (HKAS 135696, **isotype**).

Notes. Our specimens, HKAS 135697 and HKAS 135696, were grouped in a distinct clade, separated from *E. latispora* by 99% MLBS and 1.00 BIPP (Fig. 1). Our species, *E. baimana*, morphologically differs from *E. latispora* by having aseptate hairs, ascospores without oil guttules whereas the latter species possess septate hairs, asci with rounded to subconical apex, and ascospores with four or more large guttules. Therefore, *E. baimana* is introduced here as a new species.



Figure 3. *Erioscyphella baimana* (HKAS 135697, holotype) **a–c** dried ascomata on the twig **d** a vertical section of part of an ascoma **e** excipulum **f, g** hairs **h** paraphyses **i–l** asci (**k, l** asci in MLZ) **m–r** ascospores. Scale bars: 100 μ m (**d–f**); 50 μ m (**g–l**); 30 μ m (**m–r**).

***Erioscyphella gelangheica* L. Luo, K.D. Hyde, H.L. Su & C.J.Y. Li, sp. nov.**

Index Fungorum: IF902529

Facesoffungi Number: FoF16390

Fig. 4

Etymology. The epithet “*gelangheica*” refers to the collection site Gelanghe township where the holotype specimen was collected.

Holotype. HKAS135689.

Description. *Saprobic* on dead bark. **Sexual morph:** **Apothecia** scattered to partly gregarious, superficial, 0.24–0.5 mm in diameter, 0.35–0.5 mm high when dry, discoid to cupulate, long stipitate, externally covered with short, white to yellowish hairs. **Discs** concave, surface slightly rough, white to yellow. **Margin** flat to slightly involute, white to pale yellow, covered with white to pale yellow hairs. **Receptacle** discoid to cupulate, white to pale brown, clothed entirely with short, white to slightly yellow hairs. **Stipe** 0.06–0.18 mm in diameter, 0.14–0.3 mm long when dry, cylindrical, solitary, white to pale yellow, clothed with white to pale yellow hairs. **Hairs** $28\text{--}117 \times 1.6\text{--}3.9 \mu\text{m}$ ($\bar{x} = 60 \times 2.9 \mu\text{m}$, $n = 30$), clavate to cylindrical, straight to slightly curved, septate, hyaline, thin-walled, covered with hyaline granules, obtuse apex. **Hymenium** $65\text{--}120 \mu\text{m}$ ($\bar{x} = 85 \mu\text{m}$, $n = 12$), concave, surface slightly rough, light yellow in dry. **Medullary excipulum** $23.5\text{--}65 \mu\text{m}$ ($\bar{x} = 37 \mu\text{m}$, $n = 18$), thin, hyaline to light yellow, thin-walled cells of *textura intricata*, $1.3\text{--}3.6 \mu\text{m}$ ($\bar{x} = 2.2 \mu\text{m}$, $n = 50$) in diameter. **Ectal excipulum** $20\text{--}95 \mu\text{m}$ ($\bar{x} = 48 \mu\text{m}$, $n = 18$) thick, thin-walled, smooth, hyaline cells of *textura porrecta* to *textura globulosa*, $1.6\text{--}4.2 \mu\text{m}$ ($\bar{x} = 2.7 \mu\text{m}$, $n = 60$). **Paraphyses** $25\text{--}68 \times 0.9\text{--}2.0 \mu\text{m}$ ($\bar{x} = 43 \times 1.3 \mu\text{m}$, $n = 25$), longer than asci, filiform, straight, aseptate, hyaline, thin-walled, rough, with slightly acute apex. **Asci** $35\text{--}58 \times 2.2\text{--}3.60 \mu\text{m}$ ($\bar{x} = 46 \times 3.0 \mu\text{m}$, $n = 34$), 8-spored, unitunicate, clavate, straight to slightly curved, inoperculate, hyaline, wall apically thickened, laterally relatively thin, slightly smooth, with an apical, amyloid pore and tapered ends, J+ in MLZ. **Ascospores** (85/6/2) $6.0\text{--}8.3 \times 1.2\text{--}1.8 \mu\text{m}$, ($\bar{x} = 7.3 \times 1.5 \mu\text{m}$, $n = 86$), partially biserial, filiform, aseptate, thin-walled, hyaline, rough with tapering towards the obtuse ends, partially oil guttules, subspherical, hyaline, slightly smooth. **Asexual morph:** Not observed.

Material examined. CHINA • Yunnan Province, Xishuangbanna City, Menghai County, Gelanghe township, altitude 2097 m, on the decayed unidentified bark, 6 September 2022, Cuijinyi Li, LCJY1389 (HKAS 135689, **holotype**); • *ibid.*, Hongli Su, SU1978 (HKAS 135695, **paratype**).

Notes. Our specimens, HKAS 135689 and HKAS 135695, were grouped in a distinct clade, separated from the clade comprising *E. abnormis*, *E. sclerotii*, *E. ailaoensis*, *E. brasiliensis* and *E. aseptata* by 78% MLBS and 0.88 BIPP (Fig. 1). *Erioscyphella gelangheica* has shorter asci ($35\text{--}58 \mu\text{m}$ vs. $41\text{--}104 \mu\text{m}$), ascospores ($6.0\text{--}8.3 \mu\text{m}$ vs. $39\text{--}81 \mu\text{m}$), and paraphyses ($25\text{--}68 \mu\text{m}$ vs. $52\text{--}123 \mu\text{m}$) than those of *E. abnormis* (Perić and Baral 2014). *Erioscyphella gelangheica* differs from *E. sclerotii* by having long stipitate apothecia and aseptate ascospores, while *E. sclerotii* has short stipitate apothecia and 1–3-septate ascospores. Our species, *E. gelangheica*, has shorter asci ($35\text{--}58 \mu\text{m}$ vs. $85\text{--}143 \mu\text{m}$), ascospores ($6.0\text{--}8.3 \mu\text{m}$ vs. $45.5\text{--}97 \mu\text{m}$), and paraphyses ($25\text{--}68 \mu\text{m}$ vs. $100\text{--}138 \mu\text{m}$) than those of *E. ailaoensis*. Furthermore, *E. ailaoensis* has septate ascospores in contrast to the aseptate ascospores of *E. gelangheica*. Compared to long stipitate apothecia with white to pale yellow hairs and aseptate ascospores of *E. gelangheica*, *E. brasiliensis* has 0–1-septate ascospores and stipitate apothecia, with the stipe base often devoid of hairs and blue-black (Haines 1992). In addition to having aseptate ascospores, *E. gelangheica* has shorter asci ($35\text{--}58 \mu\text{m}$ vs. $70\text{--}100 \mu\text{m}$) and ascospores ($6.0\text{--}8.3 \mu\text{m}$ vs. $28.5\text{--}45.6 \mu\text{m}$) than those of *E. aseptata*, which has septate ascospores (Ekanayaka et al. 2019). Therefore, *E. gelangheica* is introduced here as a new species.



Figure 4. *Erioscyphella gelangheica* (HKAS 135689, holotype) **a–d** dried ascomata on the host **e** a vertical section of an ascoma **f** excipulum **g, h** hairs **i** paraphyses and asci **j–n** asci (**l–n** asci in MLZ) **o** apices of asci treated with Melzer's reagent **p–u** ascospores. Scale bars: 100 µm (**e**); 50 µm (**f**); 20 µm (**g–n**); 5 µm (**o–u**).

***Erioscyphella tengyueica* L. Luo, K.D. Hyde & C.J.Y. Li, sp. nov.**

Index Fungorum: IF902531

Facesoffungi Number: FoF16392

Fig. 5

Etymology. The epithet refers to the collection site of the type specimen.

Holotype. HKAS 135688.

Description. *Saprobic* on the dead twigs. **Sexual morph:** **Apothecia** superficial, scattered to partly gregarious, 0.16–0.68 mm in diameter, 0.3–0.7 mm high when dry, discoid to cupulate, shortly stipitate, externally covered with short, white hairs. **Discs** concave, surface slightly rough, white. **Margin** slightly involute, white, covered with white hairs. **Receptacle** cupulate, concolorous, clothed entirely with short, white hairs. **Stipe** 0.09–0.23 mm in diameter, 0.18–0.4 mm long when dry, cylindrical, solitary, concolorous with the receptacle, clothed with white hairs. Hairs $45\text{--}95 \times 2.8\text{--}6.9 \mu\text{m}$ ($\bar{x} = 68 \times 4.8 \mu\text{m}$, $n = 10$), clavate to cylindrical, straight or curved, septate, hyaline, thin-walled, less covered with hyaline granules, obtuse apex. **Hymenium** $65\text{--}115 \mu\text{m}$ ($\bar{x} = 87 \mu\text{m}$, $n = 12$), concave, surface slightly rough, light white in dry. **Medullary excipulum** $18\text{--}33 \mu\text{m}$ ($\bar{x} = 26 \mu\text{m}$, $n = 18$), thick, comprising hyaline, thin-walled, poorly developed cells of **textura**




Figure 5. *Erioscyphella tengyueica* (HKAS 135688, holotype) **a–c** dried ascomata on the host **d** a vertical section of an ascoma **e** excipulum **f** paraphyses **g–j** asci (**i, j** asci in MLZ) **k** ascospores. Scale bars: 100 μm (**d**); 50 μm (**e, f**); 20 μm (**g–j**); 10 μm (**k**).

globulosa, 1.4–3.5 μm ($\bar{x}$ = 2.4 μm , n = 50) in diameter. Ectal excipulum 11–30 μm ($\bar{x}$ = 21 μm , n = 18) thick, comprising thick-walled, smooth, light yellowish cells of **textura oblita** to **textura porrecta**, 1.2–3.8 μm ($\bar{x}$ = 2.2 μm , n = 60). **Paraphyses** 65–97 $\times$ 1.8–3.0 μm ($\bar{x}$ = 86 $\times$ 2.5 μm , n = 25), longer than asci, filiform, straight, aseptate, hyaline, thin-walled, narrow lanceolate, smooth, less covered with hyaline granules, with slightly obtuse apex. **Asci** 60–80 $\times$ 6.0–9.3 μm ($\bar{x}$ = 70 $\times$ 7.6 μm , n = 34), 8-spored, clavate, straight to slightly curved, inoperculate, hyaline, unitunicate, slightly smooth, with an apical, amyloid pore and rounded ends, croziers absent at the basal septum, J+ in MLZ. **Ascospores** (80/6/2) 25–31.5 $\times$ 1.6–5.5 μm , ($\bar{x}$ = 27.7 $\times$ 3.3 μm , n = 80), overlapping biserial, filiform, aseptate, thin-walled, hyaline, rough with tapering towards obtuse ends, filled with oil guttules or with 1–2 large oil guttules. **Asexual morph**: Not observed.

Material examined. CHINA • Yunnan Province, Tengchong City, Tengyue Street, altitude 1983.3 m, on the decayed unidentified twig, 21 August 2022, Cuijinyi Li, LCJY1171 (HKAS 135688, **holotype**); • *ibid.*, altitude 1774 m, on the decayed unidentified twig, 18 August 2022, Le Luo, Ly255 (HKAS 135693, **paratype**).

Notes. Our specimens, HKAS 135688 and HKAS 135693, were grouped into a distinct clade, separated from *E. papillaris* (TNS-F-81272) by 91% MLBS and 0.98 BIPP (Fig. 1). *Erioscyphella tengyueica* differs from *E. papillaris* by having aseptate ascospores and aseptate paraphyses, while the latter has septate paraphyses and aseptate or one-septate (rarely two-septate) ascospores. Our species differs from *E. otanii* by having longer asci (60–80 μm vs. 34–38.8 μm), longer ascospores (34–38.8 μm vs. 12.3–14.6 μm) and aseptate paraphyses, in contrast to the septate paraphyses of *E. otanii* (Tochihara and Hosoya 2022). Therefore, *E. tengyueica* is introduced here as a new species.

Discussion

In China, the diversity of Leotiomyces is substantial due to varied climates and ecosystems in the country (Li et al. 2022; Su et al. 2023; Guo et al. 2024; Luo et al. 2024; Zhang et al. 2024). Ongoing research continues to uncover new species and understand their roles in ecosystem functioning, highlighting the importance of preserving fungal diversity for ecological health and agricultural sustainability (Li et al. 2022; Su et al. 2023; Lestari and Chethana 2024; Luo et al. 2024).

Based on the previous research on the phylogeny, morphology and ecology of Lachnaceae (Hosoya et al. 2010), delimiting generic boundaries in Lachnaceae can no longer be defined by morphological characteristics alone (Li et al. 2022; Tochihara and Hosoya 2022; Su et al. 2023). The latest concept for *Erioscyphella* was proposed as the typical taxa characterized by the hair structures (the swollen apices, apical anamorph material/resinous material) and ascospore length (Tochihara and Hosoya 2022). This revision signifies a departure from the previously emphasized characteristics, highlighting the dynamic nature of fungal taxonomy (Tochihara and Hosoya 2022).

Despite advances in molecular techniques and the integration of morphological and ecological data, our analysis reveals persistent challenges in clarifying species boundaries. Particularly for paraphyletic members of long-spored *Lachnum* within *Erioscyphella*, the utilization of the UNITE Species Hypotheses (SH) system analysis based on ITS gene fragment, alongside traditional methods, has not provided definitive resolutions, indicating the ongoing complexity

in species delineation (Tochihara and Hosoya 2022). Phylogenetic relationships in *Erioscyphella* gradually became clear as more fresh collections were identified with sequence data (Li et al. 2022; Su et al. 2023). Almost all species of *Erioscyphella* have stable phylogenetic positions with strong statistical support, except for *E. bambusina*, which lacks sequences in public databases.

Nearly all *Erioscyphella* species occupy stable phylogenetic positions supported by strong statistical evidence, except for *E. bambusina*, which lacks sequence data in public databases. We compared the morphology of *E. bambusina* with our newly established species (Dennis 1954; Su et al. 2023). *Erioscyphella bambusina* has septate, shorter ascospores, and septate paraphyses, in contrast to the multi-septate, longer ascospores, and aseptate paraphyses of *E. ailaoensis*. *Erioscyphella gelangheica* differs from *E. bambusina* by having smaller apothecia and aseptate paraphyses, while the latter has larger apothecia and septate paraphyses. *Erioscyphella baimana* has aseptate hairs and aseptate paraphyses, while *E. tengyueica* has white discs, and J- in MLZ with and without 3% KOH pretreatment. In contrast to these two, *E. bambusina* has cream to pale yellow discs, pore blued in MLZ, and septate paraphyses.

Most records of *Erioscyphella* are from the tropics (Spooner 1987; Haines and Dumont 1983; Tochihara and Hosoya 2022). Although sample collections have extended to subtropic, temperate and cold-temperate regions (Bien and Damm 2020; Tochihara and Hosoya 2022), collected samples are still scarce. Further, it was found that the lack of available sequences of these species led to Lachnaceae and Helotiaceae taxa exhibiting paraphyletic characters in the ITS-LSU phylogeny (Johnston et al. 2019; Quandt and Haelewaters 2021). Although Johnston et al. (2019) solved this issue based on phylogenetic analyses of up to 15 concatenated genes across 279 specimens, obtaining gene sequences for multiple loci for Lachnaceae taxa is still one of the critical problems.

Continued interdisciplinary research efforts are warranted to refine our understanding of fungal taxonomy within the Lachnaceae. Future studies should explore novel methodologies, such as high-throughput sequencing and ecological niche modeling, to elucidate species boundaries and evolutionary relationships more comprehensively. Additionally, comprehensive taxonomic revisions, including detailed examination of type specimens and expanded sampling, will be crucial for resolving taxonomic ambiguities and advancing fungal systematics.

In conclusion, our study contributes to the growing body of knowledge on fungal taxonomy and highlights the need for an integrated approach combining molecular, morphological, and ecological data to address the complexities inherent in delineating generic boundaries and species relationships within the Lachnaceae family.

Acknowledgements

Le Luo thanks Dr Shaun Pennycook for his valuable assistance with the Latin binomial nomenclature.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by the Second Tibetan Plateau Scientific Expedition and Research (STEP) Program (Grant No. 2019QZKK0503) and the Chinese Research Fund (Project No. E1644111K1) under the initiative “Flexible Introduction of the High-Level Expert Program” at the Kunming Institute of Botany, Chinese Academy of Sciences. The authors extend their appreciation to the Researchers Supporting Project number (RSP2024R114), King Saud University, Riyadh, Saudi Arabia.

Author contributions

Supervision: KWTC. Writing - original draft: LL, FA. Writing - review and editing: QZ, KWTC, KDH. Collection: LL, HLS, CJYL, VL.

Author ORCIDs

Le Luo  <https://orcid.org/0009-0006-7213-0498>

Kandawatte W. Thilini Chethana  <https://orcid.org/0000-0002-5816-9269>

Qi Zhao  <https://orcid.org/0000-0001-8169-0573>

Hong-Li Su  <https://orcid.org/0000-0001-9071-7635>

Cui-Jin-Yi Li  <https://orcid.org/0000-0002-2805-7071>

Vinodhini Thiyagaraja  <https://orcid.org/0000-0002-8091-4579>

Fatimah Al-Otibi  <https://orcid.org/0000-0003-362>

Kevin D. Hyde  <https://orcid.org/0000-0002-2191-0762>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bien S, Damm U (2020) *Arboricolonus simplex* gen. et sp. nov. and novelties in *Cado-phora*, *Minutiella* and *Proliferodiscus* from Prunus wood in Germany. MycoKeys 63: 119–161. <https://doi.org/10.3897/mycokeys.63.46836>
- Cantrell SA, Haines JH (1997) New red species of *Lachnum* from the tropics. Mycological Research 101(9): 1081–1084. <https://doi.org/10.1017/S0953756297003699>
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25(15): 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Chaiwan N, Gomdola D, Wang S, Monkai J, Tibpromma S, Doilom M, Wanasinghe D, Mortimer P, Lumyong S, Hyde KD (2021) An online database providing updated information of microfungi in the Greater Mekong Subregion. Mycosphere 12(1): 1513–1526. <https://doi.org/10.5943/mycosphere/12/1/19>
- Chethana KWT, Manawasinghe IS, Hurdeal VG, Bhunjun CS, Appadoo M, Gentekaki E, Raspé O, Promputtha I, Hyde KD (2021) What are fungal species and how to delineate them? Fungal Diversity 109(1): 1–25. <https://doi.org/10.1007/s13225-021-00483-9>
- Dennis RWG (1954) Some inoperculate discomycetes of tropical America. Kew Bulletin 9(2): 289–348. <https://doi.org/10.2307/4114399>

- Ekanayaka AH, Hyde KD, Gentekaki E, McKenzie EHC, Zhao Q, Bulgakov TS, Camporesi E (2019) Preliminary classification of Leotiomyces. *Mycosphere* 10(1): 310–489. <https://doi.org/10.5943/mycosphere/10/1/7>
- Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2(2): 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Guatimosim E, Schwartsburd PB, Crous PW, Barreto RW (2016) Novel fungi from an ancient niche: Lachnoid and chalara-like fungi on ferns. *Mycological Progress* 15(12): 1239–1267. <https://doi.org/10.1007/s11557-016-1232-6>
- Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology* 59(3): 307–321. <https://doi.org/10.1093/sysbio/syq010>
- Guo MJ, Zhuo L, Wang SJ, Sui XN, Zhou H, Cai SR, Luo JT, Lei RH, Shen XY, Piepenbring M, Hou CL (2024) *Hyperdiverse Rhytismatales* on twigs of *Rhododendron* spp. *Mycosphere* 15(1): 764–880. <https://doi.org/10.5943/mycosphere/15/1/6>
- Haines JH (1992) Studies in the Hyaloscyphaceae IV: The genus *Lachnum* (Ascomycetes) of the Guayana Highlands. *Nova Hedwigia* 54(1–2): 97–112.
- Haines JH, Dumont KP (1983) Studies in the Hyaloscyphaceae II: *Proliferodiscus*, a new genus of Arachnopezizoideae. *Mycologia* 75(3): 535–543. <https://doi.org/10.1080/00275514.1983.12023717>
- Haines JH, Dumont KP (1984) Studies in the Hyaloscyphaceae III: The long-spored, lignicolous species of *Lachnum*. *Mycotaxon* 19: 1–39.
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98. https://doi.org/10.14601/phytopathol_mediterr-14998u1.29
- Han JG, An GH, Jo JW, Kim CS, Cho JH, Lee KH, Kim OT (2021) *Erioscyphella abnormis* (Lachnaceae: Ascomycota), an unrecorded species in Korea. *Journal of Asia-Pacific Biodiversity* 14(4): 662–666. <https://doi.org/10.1016/j.japb.2021.06.009>
- Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* 35(2): 518–522. <https://doi.org/10.1093/molbev/msx281>
- Hosoya T, Sasagawa R, Hosaka K, Gi-Ho S, Hirayama Y, Yamaguchi K, Toyama K, Kakishima M (2010) Molecular phylogenetic studies of *Lachnum* and its allies based on the Japanese material. *Mycoscience* 51(3): 170–181. <https://doi.org/10.1007/S10267-009-0023-1>
- Hyde KD, Noorabadi MT, Thiyagaraja V, He MQ, et al. (2024) The 2024 Outline of Fungi and fungus-like taxa. *Mycosphere* 15(1): 5146–6239.
- Index Fungorum (2024) Index Fungorum. <http://www.indexfungorum.org/names/Names.asp> [Accessed on: 2024-12-03]
- Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai YC, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu J-K, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo J-M, Ghobad-Nejhad M, Nilsson H, Pang K-L, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen T-C, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsan S, Jayawardena RS, Li W-J, Perera RH, Phookamsak R, de Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao R-L, Zhao Q, Kang J-C, Prompttha I (2015) The Faces of Fungi database: Fungal names

- linked with morphology, phylogeny and human impacts. *Fungal Diversity* 74(1): 3–18. <https://doi.org/10.1007/s13225-015-0351-8>
- Johnson M, Zaretskaya I, Raytselis Y, Merezuk Y, McGinnis S, Madden TL (2008) NCBI BLAST: A better web interface. *Nucleic Acids Research* 36(Web Server): W5–W9. <https://doi.org/10.1093/nar/gkn201>
- Johnston PR, Quijada L, Smith CA, Baral H-O, Hosoya T, Baschien C, Pärtel K, Zhuang W-Y, Haelewaters D, Park D, Carl S, López-Giráldez F, Wang Z, Townsend JP (2019) A multigene phylogeny toward a new phylogenetic classification of Leotiomyces. *IMA Fungus* 10(1): 1–22. <https://doi.org/10.1186/s43008-019-0002-x>
- Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kirschstein W (1939) Über neue, seltene und kritische Ascomyceten und Fungi imperfecti. II. *Annales Mycologici* 37: 88–140.
- Korf RP (1978) Nomenclatural and taxonomic notes on *Lasiobelonium*, *Erioscypha* and *Erioscyphella*. *Mycotaxon* 7: 399–406.
- Lestari A, Chethana KWT (2024) *Lambertella* (Rutstroemiaceae, Helotiales) from Northern Thailand. *New Zealand Journal of Botany* 1: 62(2–3): 443–480. <https://doi.org/10.1080/0028825X.2024.2344859>
- Li CJY, Chethana KWT, Lu ZY, Zhao Q (2022) Two novel species of Lachnaceae (Helotiales, Leotiomyces) from southwestern China. *Current Research in Environmental & Applied Mycology (Journal of Fungal Biology)* 12: 333–345. <https://doi.org/10.5943/cream/12/1/20>
- Li CJY, Thilini Chethana KW, Eungwanichayapant PD, Zhou DQ, Zhao Q (2024a) Additional four species of *Tatraea* (Leotiomyces, Helotiales) in Yunnan Province, China. *MycoKeys* 102: 127–154. <https://doi.org/10.3897/mycokeys.102.112565>
- Li CJY, Chethana KWT, Eungwanichayapant PD, Luo L, Yang ZL, Dong WJ, Guo YY, Liu CH, Alotibi F, Hyde KD, Zhao Q (2024b) *Longistipes* gen. nov. and four novel species of Hyphodiscaceae along with six new collections of Leotiomyces in Yunnan Province, China. *Mycosphere* 15(1): 4744–4787. <https://doi.org/10.5943/mycosphere/15/1/20>
- Liu YJ, Whelen S, Hall BD (1999) Phylogenetic relationships among ascomycetes: Evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16(12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Luo L, Wei DP, Zhao Q, Chethana KWT (2024) Unveiling the Diversity: A novel species of *Dicephalospora* (Helotiaceae, Helotiales) discovered in China. *Phytotaxa* 652: 59–68. <https://doi.org/10.11646/phytotaxa.652.1.5>
- Maharachchikumbura SSN, Chen Y, Ariyawansa HA, Hyde KD, Haelewaters D, Perera RH, Samarakoon MC, Wanasinghe DN, Bustamante DE, Liu J, Lawrence DP, Cheewangkoon R, Stadler M (2021) Integrative approaches for species delimitation in Ascomycota. *Fungal Diversity* 109(1): 155–179. <https://doi.org/10.1007/s13225-021-00486-6>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, 1–8. [14 Nov. 2010] <https://doi.org/10.1109/GCE.2010.5676129>

- Nagao H (1996) Discomycetes on decayed tree fern. (2) *Lachnum varians* (Rehm) Spooner and *Lachnum sclerotii* (AL Smith) Haines et Dumont new to Japan. Bulletin of the National Science Museum 22: 105–111.
- Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Niego AGT, Rapior S, Thongklang N, Raspé O, Hyde KD, Mortimer P (2023a) Reviewing the contributions of macrofungi to forest ecosystem processes and services. Fungal Biology Reviews 44: 100294. <https://doi.org/10.1016/j.fbr.2022.11.002>
- Niego AGT, Lambert C, Mortimer P, Thongklang N, Rapior S, Grosse M, Schrey H, Charria-Girón E, Walker A, Hyde KD, Stadler M (2023b) The contribution of fungi to the global economy. Fungal Diversity 121(1): 95–137. <https://doi.org/10.1007/s13225-023-00520-9>
- Perić B, Baral HO (2014) *Erioscyphella curvispora* spec. nov. from Montenegro. Mycologia Montenegrina 17: 89–104.
- Quandt CA, Haelewaters D (2021) Phylogenetic advances in Leotiomyces, an understudied clade of taxonomically and ecologically diverse fungi. Encyclopedia of Mycology 1: 284–294. <https://doi.org/10.1016/B978-0-12-819990-9.00052-4>
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Spooner BM (1987) Helotiales. of Australasia: Geoglossaceae, Orbiliaceae, Sclerotiniaceae, Hyaloscyphaceae. Bibliotheca Mycologica 116: 470–474.
- Su HL, Chethana KWT, Li L, Li W, Zhao Q (2022) *Diplocarpa constans* sp. nov., a new species of Cordieritidaceae from Yunnan, China. Phytotaxa 561(1): 75–84. <https://doi.org/10.11646/phytotaxa.561.1.7>
- Su HL, Chethana KWT, Zeng M, Zhao Q (2023) Two new species of *Erioscyphella* (Lachnaceae) from southwestern. Current Research in Environmental & Applied Mycology 13(1): 16–33. <https://doi.org/10.5943/cream/13/1/2> [Journal of Fungal Biology]
- Tello S, Baral HO (2016) *Erioscyphella lunata* (Lachnaceae), a rare discomycete collected in Spain. Ascomycete.Org : Revue Internationale pour la Taxinomie des Ascomycota 8(4): 157–162. <https://doi.org/10.25664/ART-0183>
- Thiyagaraja V, He SC, Luo L, Meng Q, Yang HD, Yang YY (2024) Research profiles – I: The research of Vinodhini Thiyagaraja and her team at Kunming Institute of Botany, CAS, P. R. China. Current Research in Environmental & Applied Mycology 14(1): 49–62. <https://doi.org/10.5943/cream/14/1/2>
- Tochihara Y, Hosoya T (2019) Three new species of *Incrucipulum* (Lachnaceae, Helotiales, Ascomycota) from Japan. Phytotaxa 403(1): 25–38. <https://doi.org/10.11646/phytotaxa.403.1.2>
- Tochihara Y, Hosoya T (2022) Examination of the generic concept and species boundaries of the genus *Erioscyphella* (Lachnaceae, Helotiales, Ascomycota) with the proposal of new species and new combinations based on the Japanese materials. MycoKeys 87: 1–52. <https://doi.org/10.3897/mycokeys.87.73082>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several Cryptococcus species. Journal of Bacteriology 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>

- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications 18: 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao R, Aptroot A, Leontyev D, Saxena R (2020) Outline of Fungi and fungus-like taxa. Mycosphere 11(1): 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>
- Wijayawardene NN, Hyde K, Dai DQ, Sánchez-García M, Goto BT, Saxena RK, Erdoğan M, Selçuk F, Rajeshkumar KC, Aptroot A, Błaszczowski J, Boonyuen N, da Silva GA, de Souza FA, Dong W, Ertz D, Haelewaters D, Jones EBG, Karunarathna SC, Kirk PM, Kukwa M, Kumla J, Leontyev DV, Lumbsch HT, Maharachchikumbura SSN, Marguno F, Martínez-Rodríguez P, Mešić A, Monteiro JS, Oehl F, Pawłowska J, Pem D, Pfliegler WP, Phillips AJL, Pošta A, He MQ, Li JX, Raza M, Sruthi OP, Suetrong S, Suwannarach N, Tedersoo L, Thiyagaraja V, Tibpromma S, Tkálčec Z, Tokarev YS, Wanasinghe DN, Wijesundara DSA, Wimalaseana SDMK, Madrid H, Zhang GQ, Gao Y, Sánchez-Castro I, Tang LZ, Stadler M, Yurkov A, Thines M (2022) Outline of Fungi and fungus-like taxa – 2021. Mycosphere 13(1): 53–453. <https://doi.org/10.5943/mycosphere/13/1/2>
- Zeng XY, Tan TJ, Tian FH, Wang Y, Wen TC (2023) OFPT: A one-stop software for fungal phylogeny. Mycosphere 14(1): 1730–1741. <https://doi.org/10.5943/mycosphere/14/1/20>
- Zhao P, Zhuang WY (2011) Evaluation of ITS region as a possible DNA barcode for the genus Lachnum (Helotiales). Mycosystema 30(6): 932–937.
- Zhang ZY, Pan H, Tao G, Li X, Han YF, Feng Y, Tong SQ, Ding CY (2024) Culturable mycobiota from Guizhou Wildlife Park in China. Mycosphere 15(1): 654–763. <https://doi.org/10.5943/mycosphere/15/1/5>
- Zoller S, Scheidegger C, Sperisen C (1999) PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. Lichenologist 31(5): 511–516. <https://doi.org/10.1006/lich.1999.0220>